

M	T	W	T	F	S	S
Page No.					YOUVA	Date:

216121

Class \Rightarrow B.Sc. Part I Hons.

Subject \Rightarrow Chemistry

Paper \Rightarrow IA (Physical chemistry)

Chapter \Rightarrow Gaseous state (Group-A)

Topic \Rightarrow Different type of velocities

Name \Rightarrow Dr. Amarendra Kumar

Dept. of chemistry

Jain College, Ara.

Most Probable velocity

The velocity possessed by maximum number of molecules of a gas at a given temperature is called most probable velocity.

According to the calculations made by Maxwell, the most probable velocity (V_{mp}) is given by the expression

$$V_{mp} = \sqrt{\frac{2RT}{M}}$$

Substituting the values of R, T and M in this expression, the most probable velocity can be calculated.

Average velocity

The velocity possessed by maximum number of molecules of a gas at a given temperature is called Average velocity.

Let n molecules of a gas having individual velocities $v_1, v_2, v_3, \dots, v_n$

$$\therefore \text{Average velocity} = \frac{v_1 + v_2 + v_3 + \dots + v_n}{n}$$

from Maxwell equation it has been established that the average velocity v_{av} is given by the expression

$$v_{av} = \sqrt{\frac{8RT}{\pi M}}$$

Substituting the values of R, T, π and M , in this expression, the average velocity can be calculated.

Root Mean Square Velocity

The square root of the mean of the squares of different velocities possessed by molecules of a gas at a given temperature is called root mean square velocity.

It is denoted by u .

If $v_1, v_2, v_3, \dots, v_n$ are the velocities of n molecules in a gas. Then if the mean of the squares of all the velocities is

$$u^2 = \frac{v_1^2 + v_2^2 + v_3^2 + \dots + v_n^2}{n}$$

Taking the root

$$u = \sqrt{\frac{v_1^2 + v_2^2 + v_3^2 + \dots + v_n^2}{n}}$$

u is the root mean square velocity or RMS velocity.

The value of root mean square velocity u at a given temperature is calculated from the kinetic gas equation.

(3)

M	T	W	T	F	S	S
Page No.:		Date:		YOUVA		

$$PV = \frac{1}{3} m N u^2$$

$$\text{or } u^2 = \frac{3PV}{mN}$$

for 1 mole of ggs

$$PV = RT$$

$$\therefore u^2 = \frac{3RT}{M}$$

Where M = Molar Mass

$$\text{or } u = \sqrt{\frac{3RT}{M}}$$

Substituting the values of R , T and M , the Root mean square velocity u can be calculated.

Relation between Average velocity, Root mean square velocity and Most probable velocity

We know that,

$$\text{Average velocity } (\bar{v}) = \sqrt{\frac{8RT}{\pi M}}$$

$$\text{Root mean square velocity } u = \sqrt{\frac{3RT}{M}}$$

$$\therefore \frac{\bar{v}}{u} = \sqrt{\frac{8RT}{\pi M}} \times \sqrt{\frac{M}{3RT}}$$

$$= \frac{8}{\sqrt{3\pi}}$$

$$= 0.9213$$

$$\text{Or } \bar{v} = u \times 0.9213$$

\therefore Average velocity = $0.9213 \times$ Room mean square velocity

(4)

M	T	W	T	F	S	S
Page No.:	YOUVA					
Date:						

The most probable velocity.

$$V_{mp} = \sqrt{\frac{2RT}{M}}$$

$$\text{and } u = \sqrt{\frac{3RT}{M}}$$

$$\therefore \frac{V_{mp}}{u} = \frac{\sqrt{\frac{2RT}{M}}}{\sqrt{\frac{3RT}{M}}} = \frac{\sqrt{2/3}}{\sqrt{3}} = \frac{1}{\sqrt{3}}$$

$$= \frac{1}{\sqrt{3}} = 0.8165$$

Most probable velocity = $0.8165 \times \text{Root mean square}$

Root mean square velocity is calculated by the application of kinetic gas equation.

Knowing the value of RMS velocity we can calculate the average velocity and most probable velocity.