

class  $\Rightarrow$  B.Sc.(Part I)'s subsidiary  
subject  $\Rightarrow$  chemistry

chapter  $\Rightarrow$  Gaseous state (Grp.A)

Topic  $\Rightarrow$  Real and Ideal gases,  
Deviation of Real gases from ideal behaviour.

Name  $\Rightarrow$  Dr. Amarendra Kumar

Dept. of chemistry -

Jain College, Ara.

### Real gases

A real gas is one which obeys the gas laws fairly well under low pressure or high temperature.

All gases are real gases. They show more and more deviations from the gas laws as the pressure is increased or the temperature is decreased.

### Ideal gas

An ideal gas is one which obeys the gas laws or the gas equation ( $PV = nRT$ ) at all pressure and temperature.

However no gas is ideal. Almost all gases show significant deviations from the ideal behaviour.

### Difference between Ideal gas and

#### Real gases

(1) An ideal gas obeys the basic laws strictly under all conditions of temperature and pressure.

Real gases do not obey the basic laws under all conditions of temperature and pressure.

(2)

(2) In an ideal gas the actual volume of molecules is negligible.

While in Real gases the actual volume of molecules is appreciable.

(3) In Ideal g<sub>as</sub> there are no attractive forces between molecules.

While in Real g<sub>ases</sub> attractive forces between molecules exist.

(4) Molecular collisions in an ideal g<sub>as</sub> are perfectly elastic while it is not so in a real g<sub>as</sub>.


### Deviation of Real gases from ideal behaviour

According to Boyle's law,  $PV = \text{constant}$  at constant temp. This means that the product  $PV$  should remain constant at all pressures provided temperature is kept constant.

Thus a plot of  $PV$  vs  $P$  should be a horizontal straight line parallel to  $x$ -axis.

However, in actual practice, it is found that no gas gives a straight line plot. Two types of curves are generally obtained.

fig:- Plots of  $PV$  vs  $P$  for different gases at  $0^\circ\text{C}$  ( $273\text{K}$ )



M	T	W	T	F	S	S
Page No.:	YOUVA					
Date:						

(3)

(i) For gases like hydrogen and helium, the product  $PV$  continuously increases with increase of pressure. (from plot  $PV$  vs  $P$ )

(ii) For gases like  $\text{CO}$  and  $\text{CH}_4$ , the product  $PV$  first decreases with increase of pressure, reaches a minimum value and then begins to increase (from plot  $PV$  vs  $P$ ).   
 —X—

→ Deviation of real gases from ideal behaviour in terms of compressibility factor, denoted by  $Z$ .

$$Z = \frac{PV}{nRT}$$

for an Ideal gas,  $Z=1$  at all temperature and pressure.

In case of real gases the compressibility factor  $Z$  varies from values less than 1 to values greater than 1 with changes of temperature and pressure.

—X—