



Class ⇒ B.Sc. (Part-II) subsidiary
 subject ⇒ chemistry
 Chapter ⇒ solid state
 Topic ⇒ Lattice energy and
 Born-Haber cycle

Name ⇒ Dr. Amarendra Kumar
 Deptt. of chemistry
 Jain College, Ara.

Lattice Energy of an ionic crystal (Born-Haber cycle)

The amount of energy released when cations and anions in their gaseous state are brought together from infinite separation to form a crystal is called lattice Energy.



Experimentally, the lattice enthalpy of an ionic compound is determined by using the Born-Haber cycle.




fig:- A Born-Haber cycle for the formation of NaCl crystal from its elements.


The positive and negative ion in an ionic crystal are held together by electrostatic forces.

(2)

Determination of Lattice Energy


The formation of an ionic solid from its elements can be thought of as occurring directly in one step or indirectly as the sum of the series of steps.

Let us illustrate by taking example of NaCl. Enthalpy change for direct formation \Rightarrow The enthalpy change for the direct formation of sodium chloride from sodium metal and chlorine is -411 kJ .




Enthalpy change by indirect steps \Rightarrow The direct enthalpy change of NaCl is equal to enthalpy changes of five steps by which the formation of NaCl can be accomplished.


Step 1 \Rightarrow Conversion of sodium metal to gaseous atoms (sublimation).




Step 2 \Rightarrow Dissociation of chlorine molecules to chlorine atoms. The enthalpy of dissociation is 121 kJ per mole of chlorine.



Step 3 \Rightarrow Conversion of gaseous atom to sodium ions by losing electrons. The enthalpy of ionisation is the ionisation energy, 495 kJ/mole of sodium atom.



Step 4 \Rightarrow Chlorine atoms gain an electron to form chloride ions. The energy released is the electron affinity of chlorine -348 kJ




Step 5 \Rightarrow Sodium and chloride ions get together

M	T	W	T	F	S	S
Page No.:	YOUVA					
Date:						

(3)

and form the crystal lattice. The energy released in this process is the negative of lattice energy.



The lattice energy can be calculated by equating the enthalpy of formation of NaCl, -411 kJ, to the sum of the enthalpy changes for five steps.

$$\Delta H_1^\circ + \Delta H_2^\circ + \Delta H_3^\circ + \Delta H_4^\circ + \Delta H_5^\circ = -411 \text{ kJ}$$

$$108 \text{ kJ} + 121 \text{ kJ} + 495 \text{ kJ} - 348 \text{ kJ} - \text{lattice energy} = -411 \text{ kJ}$$

By solving this equation, we get

$$\text{Lattice energy} = +787 \text{ kJ mol}^{-1}$$

The cycle of changes shown in figure is called the Born-Haber cycle.