

B.Sc Part I (Hons.)

Fundamental theorem of Algebra. Problems

Prove that in equation with real coefficients imaginary roots occur in conjugate pairs.

Proof: — Let  $\alpha+i\beta$  be the root of the equation  $f(x)=0$  in which the coefficients are real, then  $\alpha-i\beta$  must also be a root of the equation  $f(x)=0$ .

When  $\alpha+i\beta$  is a root of  $f(x)=0$  then

$$f(\alpha+i\beta)=0 \quad \text{--- (1)}$$

Now to follow division algorithm we divide the polynomial  $f(x)$  by

$$[(x-(\alpha+i\beta)) [x-(\alpha-i\beta)] \text{ i.e. by } (x-\alpha)^2+\beta^2.$$

If the quotient be  $Q$  and remainder be  $Rx+R'$  then we have

$$f(x) = \{(x-\alpha)^2+\beta^2\} Q + (Rx+R') \quad \text{--- (i)}$$

As  $\alpha+i\beta$  is a root of equation then

$$f(\alpha+i\beta)=0, \text{ then by (1)}$$

$$f(\alpha+i\beta)=0 = \{(\alpha+i\beta)-\alpha\}^2+\beta^2\} Q + R(\alpha+i\beta)+R'$$

$$\Rightarrow 0 = R(\alpha+i\beta)+R'$$

$$\Rightarrow R\alpha+R'+Ri\beta=0+0$$

Equating real and imaginary part, we get

$$R\alpha+R'=0 \text{ and } R\beta=0, \text{ i.e. } R=0 \text{ and } \beta=0$$

$$\therefore 0\alpha+R'=0 \Rightarrow R'=0$$

We have  $R=0$  and  $R'=0$ , so by (i)

$$f(x) = [(x-\alpha)^2 + \beta^2]Q + 0 = (x-\alpha)^2 + \beta^2$$

$$= [(\alpha-x)^2 + (\beta i)^2]Q$$

$$f(x) = (x-\alpha-i\beta)(x-\alpha+i\beta)Q$$

This shows that  $\alpha - \alpha - i\beta$  is a factor of  $f(x)$ . i.e.  $x = \alpha + i\beta$  is a root of  $f(x) = 0$ .

That Proved.

Question:- From the equation with rational coefficients which shall have for two of its roots are  $\sqrt{3}$  and  $2+i$ .

Ans:- We have when one of the roots of given equation is  $\sqrt{3}$  then other must be  $-\sqrt{3}$  and when one root is  $2+i$  then other will be  $2-i$ .

The required equation will be

$$\{x-(2+i)\} \{x-(2-i)\} \{(x-\sqrt{3})(x+\sqrt{3})\} = 0$$

$$\{(x-2)-i\} \{(x-2)+i\} \{x^2 - (\sqrt{3})^2\} = 0$$

$$\Rightarrow \{(x-2)^2 - i^2\} (x^2 - 3) = 0$$

$$\{(x-2)^2 + 1\} (x^2 - 3) = 0$$

$$(x^2 + 4 - 4x + 1) (x^2 - 3) = 0$$

$$(x^2 + 4x + x^2 - 4x + 5) (x^2 - 3) = 0$$

$$x^4 - 4x^3 + 2x^2 + 12x + 5 = 0$$

===== Arjani Kumar Singh =====