

PROTOTHERIA

AND

METATHERIA


## PROTOHEREA & METATHERIA.

INTRODUCTION:- Mammals, the youngest baby of the animal family has overpowered the whole physical world by development of its brain since the time immemorial. Some of the mammals are deleterious, some are domesticated due to their helpful behaviour. But the infatuation of human being himself a mammal has crumble due to the nonsense behaviours of certain mammals. All these mammals from deleterious to the domesticated are grouped under viviparous (mammals laying the young ones) but the Keenast mind of embryologist claimed that some mammals are oviparous (egg laying Mammals). All the mammals ranging from oviparous to the viviparous residing in the water, in the forests, and on the ground are broadly grouped into three prototherians and eutherians.


### STRUCTURE PECULIARITIES OF PROTOHEREA:-

#### External features:-


- ① Body small and thickly covered with hairs, which are interspersed with strong pointed spines.
- ② The jaws form a long pointed snout or Rostrum which is tactile in nature in case of echidna but in Ornithorhynchus the upper jaw forms a broad flat muzzle like beak of a duck.



- ④ You have no external fins.
- ⑤ You are small with no webbed tail membrane in case of Onithorhynchus the tail is short and broad.



- ⑥ In case of Onithorhynchus beyond the claws of forelimb is a web having leathery extension the web is used for swimming.
- ⑦ In case of hind limb of Echidna the second digit has a long curved talon-claw to clean the spines and hairs. In Onithorhynchus the foot has a smaller web below the first digit the web forms a caudal pointed prolongation.
- ⑧ In the hind limb of male Echidna is a horny spur on the inner side of tarsus and a duct from poison gland in the thigh opens into the spur. It is more developed in Onithorhynchus than Echidna.



Onithorhynchus ← post leg → Echidna

mammary glands are without nipples. The male also has mammary glands and secretes milk, this condition is known as gynecomastism in which both parents share in feeding the young.


Body Cavity: The body cavity is divided into the dorsal thoracic cavity and the ventral abdominal cavity by a horizontal muscular partition, the diaphragm.

### Endoskeleton:

- ① The skull is dicnoidal and structures between the bones are not distinct.
- ② The sphenoid bone is absent but ectopterygoid bone is present.
- ③ The tympanic bulla is absent and the malleus ossicles are comparatively larger than usual.
- ④ Each mandible is formed of a single dentary bone.
- ⑤ The vertebrae are without epiphysis and the ribs are single headed.
- ⑥ T shaped interclavicle is present.
- ⑦ The pelvic girdle has an additional epipubic bone. Acetabulum is perforated and the ischia and pubis are fused at a ventral symphysis.

### Digestive System:

- ① Teeth are absent at any stage in echidna but in ornithorhynchus the young ones bears the teeth and the adults possess epidermal plates which assist in the process of mastication.
- ② Muzzle is used for capturing worms and molluscs from mud of river beds which can be stored in cheek pouches. Tongue is long.



- ⑩ Dorsal appendages to store their faecal :  
In leeches there is a long proboscis tongue and large salivary gland. The posterior surface of the tongue has horny serrations which grind the insects. The saliva neutralizes the formic acid of the ants.

### Circulatory System:

- ① The heart is four chambered but the atriaco ventricular valve is incomplete and muscular.
- ② choroid - tendines are absent.
- ③ only left aortic arch persists in the adult.
- ④ The RBC is non-nucleated.

### Nervous System:

- ① Brain is poorly developed.
- ② The corpus callosum is absent but the anterior commissure is large ③ The cochlea is less coiled.

### Urogenital system:-

- ① The kidneys are metanephric and without open into urogenital sinus which does not traverse the penis.
- ② The testes are abdominal.
- ③ Penis consists of a spongy Corpus spongiosum and corpus fibrosum and bears a groove for transmitting spermatozoa but not the urine.
- ④ Right ovary is reduced.
- ⑤ The oviducts open separately into the cloaca.
- ⑥ The vagina and uterus are absent.

## STRUCTURE PECULIARITIES OF METATHERIA

### External features:-

- ① Body is enclosed in a furry integument.
- ② The pinna is present.
- ③ Tail is long and prehensile & acts as a balancing while running and also takes a firm hold of twigs etc.
- ④ Locomotion is bipedal with modifications in ilia and thigh muscles the Kangaroo moves rapidly by long, erect springs of over 25 ft. by jumping, the tail is used for balancing.
- ⑤ Hind limbs are long and powerful, the foot has four digits, the hallux being absent. Second and third metatarsals and digits are thin and small, these two digits are united together by integument so that the foot appears three-toed. The fourth metatarsals and digit are very large with a strong claw which is used in fighting the fifth toe is small.
- ⑥ In female the abdomen possesses a sac like structure the marsupium which carries under developed young.
- ⑦ Marsupium encloses the nipples of the mammary glands to feed the young ones.

### Endoskeleton:-

- ① The skull is dicnoidal and the structures are distinct. Alisphenoid bone is present.
- ② Tympanic bulla is formed by alisphenoid bone.
- ③ Vertebrae have prominent and there are no cervical ribs sacrum is formed of a single vertebra.
- ④ Pectoral girdle has large scapula with a spine. Coracoid is reduced. Clavicles are large but there is no interclavicle.
- ⑤ Epipubic bones in the pelvic girdle are present.
- ⑥ Ischium and pubis are fused at the ventral synostosis.

## Digestive System:-

- ① Teeth are peculiar in being monophyodont and heterodont.
- ② Teeth are numerous in number with 9 incisor in each upper jaw and 3 in the lower jaw, there are 3 premolars, and 4 molars in each half of the each jaw molars have grinding surfaces.

## Circulatory System:-

- ① The heart is completely divided into four chambers and the auriculo-ventricular valve is membranous.
- ② Each superior vena cava receives an azygous vein.

## Nervous System:-

- ① The olfactory lobes are comparatively larger.
- ② The cerebral hemispheres are small and do not extend posteriorly over the cerebellum.
- ③ A corpus callosum is absent, but the anterior commissure is well developed.
- ④ The cochlea of internal ear is very much coiled.

## Urino-genital System:-

- ① The kidneys are metanephric.
- ② The both sexes are water pass between the genital ducts.
- ③ The testes are extra abdominal and lies in the scrotal sac in front of the penis.
- ④ The female possesses two vaginal and two uteri which one is primitive? - Having a birds eye view on the above mentioned account regarding the structural peculiarities of prototheria and metatheria it can be undoubtedly said that prototheria is more primitive than the metatheria on the basis of the following characters:-

- ① Hair on the body of echidna traversed by the pores.
- ② Spines
- ③ poison spur present on the hind leg.
- ④ pinnae as the external ear are absent.
- ⑤ structures in the skull bone are indistinct.
- ⑥ tympanic bulla is absent.
- ⑦ vertebrae devoid of epiphyses.
- ⑧ absence of uterus and vagina in the female.
- ⑨ eggs laying in nature.
- ⑩ absence of nipples in the mammary glands.
- ⑪ cloaca is present.
- ⑫ Atriculo ventricle valve is incomplete and muscular.

Affinity with Prototheria - The developed characters having much similarities with Prototheria thus are given below.

- ① cloaca is found.
- ② corpus callosum is absent.
- ③ large olfactory lobes are found.
- ④ epipubic bones are found in girdle.
- ⑤ clavicle is found in pectoral girdle.
- ⑥ Similarities with Eutheria
- ① There are 10 pairs; egg laying in nature.
- ② Nipples found in mammary gland.
- ③ yolk is absent.
- ④ cleavage is of holoblastic type.
- ⑤ Pinna present.
- ⑥ heterodont teeth are found.
- ⑦ placenta absent.
- ⑧ Placenta are found.
- ⑨ Uterus & vagina is also found.